

BROVANA® (arformoterol tartrate) Inhalation Solution 15 mcg * /2 mL
*potency expressed as arformoterol
For oral inhalation only
WARNING: ASTHMA RELATED DEATH Long-acting beta₂-adrenergic agonists (LABA) increase the risk of asthma-related death. Data from a large placebo-controlled US study that compared the safety of another long-acting beta₂-adrenergic agonist (salmeterol) or placebo added to usual asthma therapy showed an increase in asthma-related deaths in patients receiving salmeterol. This finding with salmeterol is considered a class effect of LABA, including arformoterol, the active ingredient in BROVANA (see WARNINGS). The safety and efficacy of BROVANA in patients with asthma have not been established. All LABA, including BROVANA, are contraindicated in patients with asthma without use of a long-term asthma control medication (see CONTRAINDICATIONS).
DESCRIPTION BROVANA (arformoterol tartrate) Inhalation Solution is a sterile, clear, colorless, aqueous solution of the tartrate salt of arformoterol, the (R,R)-enantiomer of formoterol. Arformoterol is a selective beta ₂ -adrenergic bronchodilator. The chemical name for arformoterol tartrate is formamide, N-[2-hydroxy-5-[(1R)-1-hydroxy-2-[[[(1R)-2-(4-methoxyphenyl)-1-methylethyl] amino]ethyl] phenyl]-, (2R,3R)-2,3-dihydroxybutanedioate (1:1 salt), and its established structural formula is as follows: <div style="display: flex; justify-content: space-around; align-items: center;"><div style="text-align: center;"><p>Chemical structure of arformoterol tartrate, showing the (R,R)-enantiomer of formoterol tartrate.</p></div><div style="text-align: center;"><p>Chemical structure of tartaric acid, the counterion of the tartrate salt.</p></div></div>
The molecular weight of <i>arformoterol tartrate</i> is 494.5 g/mol, and its empirical formula is C ₁₉ H ₂₄ N ₂ O ₄ • C ₄ H ₆ O ₆ (1:1 salt). It is a white to off-white solid that is slightly soluble in water. Arformoterol tartrate is the United States Adopted Name (USAN) for (R,R)-formoterol L-tartrate. BROVANA is supplied as 2 mL of arformoterol tartrate solution packaged in 2.1 mL unit-dose, low-density polyethylene (LDPE) ready-to-use vials. Each ready-to-use vial contains 15 mcg of arformoterol (equivalent to 22 mcg of arformoterol tartrate) in a sterile, isotonic saline solution, pH-adjusted to 5.0 with citric acid and sodium citrate. BROVANA requires no dilution before administration by nebulization. Like all other nebulized treatments, the amount delivered to the lungs will depend upon patient factors, the nebulizer used, and compressor performance. Using the PARI LC PLUS® nebulizer (with mouthpiece) connected to a PARI DUARA-NEB® 3000 compressor under <i>in vitro</i> conditions, the mean delivered dose from the mouthpiece (% nominal) was approximately 4.1 mcg (27.6%) at a mean flow rate of 3.3 L/min. The mean nebulization time was 6 minutes or less. BROVANA should be administered from a standard jet nebulizer at adequate flow rates via face mask or mouthpiece (see Dosage and Administration). Patients should be carefully instructed on the correct use of this drug product (please refer to the accompanying Medication Guide).
CLINICAL PHARMACOLOGY Mechanism of Action Arformoterol, the (R,R)-enantiomer of formoterol, is a selective long-acting beta ₂ -adrenergic receptor agonist (beta ₂ -agonist) that has two-fold greater potency than racemic formoterol (which contains both the (S,S) and (R,R)-enantiomers). The (S,S)-enantiomer is about 1,000-fold less potent as a beta ₂ -agonist than the (R,R)-enantiomer. While it is recognized that beta ₂ -receptors are the predominant adrenergic receptors in bronchial smooth muscle and beta ₂ -receptors are the predominant receptors in the heart, data indicate that there are also beta ₂ -receptors in the human heart comprising 10% to 50% of the total beta-adrenergic receptors. The precise function of these receptors has not been established, but they raise the possibility that even highly selective beta ₂ -agonists may have cardiac effects. The pharmacologic effects of beta ₂ -adrenoceptor agonist drugs, including arformoterol, are at least in part attributable to stimulation of intracellular adenylyl cyclase, the enzyme that catalyzes the conversion of adenosine triphosphate (ATP) to cyclic-3',5'-adenosine monophosphate (cyclic AMP). Increased intracellular cyclic AMP levels cause relaxation of bronchial smooth muscle and inhibition of release of mediators of immediate hypersensitivity from cells, especially from mast cells. <i>In vitro</i> tests show that arformoterol is an inhibitor of the release of mast cell mediators, such as histamine and leukotrienes, from the human lung. Arformoterol also inhibits histamine-induced plasma albumin extravasation in anesthetized guinea pigs and inhibits allergen-induced eosinophil influx in dogs with airway hyper-responsiveness. The relevance of these <i>in vitro</i> and animal findings to humans is unknown. Animal Pharmacology In animal studies investigating its cardiovascular effects, arformoterol induced dose-dependent increases in heart rate and decreases in blood pressure consistent with its pharmacology as a beta-adrenergic agonist. In dogs, at systemic exposures higher than anticipated clinically, arformoterol also induced exaggerated pharmacologic effects of a beta-adrenergic agonist on cardiac function as measured by electrocardiogram (sinus tachycardia, atrial premature beats, ventricular escape beats, PVCs). Studies in laboratory animals (minipigs, rodents, and dogs) have demonstrated the occurrence of arrhythmias and sudden death (with histologic evidence of myocardial necrosis) when beta-agonists and methylxanthines are administered concurrently. The clinical significance of these findings is unknown.
Pharmacokinetics The pharmacokinetics (PK) of arformoterol have been investigated in healthy subjects, elderly subjects, renally and hepatically impaired subjects, and chronic obstructive pulmonary disease (COPD) patients following the nebulization of the recommended therapeutic dose and doses up to 96 mcg. Absorption In COPD patients administered 15 mcg arformoterol every 12 hours for 14 days, the mean steady-state peak (R,R)-formoterol plasma concentration (C _{max}) and systemic exposure (AUC _{0-12h}) were 4.3 pg/mL and 34.5 pg•hr/mL, respectively. The median steady-state peak (R,R)-formoterol plasma concentration time (t _{max}) was observed approximately one half hour after drug administration. Systemic exposure to (R,R)-formoterol increased linearly with dose in COPD patients following arformoterol doses of 5 mcg, 15 mcg, or 25 mcg twice daily for 2 weeks or 15 mcg, 25 mcg, or 50 mcg once daily for 2 weeks. In a crossover study in patients with COPD, when arformoterol 15 mcg inhalation solution and 12 and 24 mcg formoterol fumarate inhalation powder (Foradil® Aerolizer™) was administered twice daily for 2 weeks, the accumulation index was approximately 2.5 based on the plasma (R,R)-formoterol concentrations in all three treatments. At steady state, geometric means of systemic exposure (AUC _{0-12h}) to (R,R)-formoterol following 15 mcg of arformoterol inhalation solution and 12 mcg of formoterol fumarate inhalation powder were 39.33 pg•hr/mL and 33.93 pg•hr/mL, respectively (ratio 1.16; 90% CI 1.00, 1.35), while the geometric means of the C _{max} were 4.30 pg/mL and 4.75 pg/mL, respectively (ratio 0.91; 90% CI 0.76, 1.09). In a study in patients with asthma, treatment with arformoterol 50 mcg with pre- and post-treatment with activated charcoal resulted in a geometric mean decrease in (R,R)-formoterol AUC _{0-6h} by 27% and C _{max} by 23% as compared to treatment with arformoterol 50 mcg alone. This suggests that a substantial portion of systemic drug exposure is due to pulmonary absorption. Distribution The binding of arformoterol to human plasma proteins <i>in vitro</i> was 52-65% at concentrations of 0.25, 0.5 and 1.0 ng/mL of radiolabeled arformoterol. The concentrations of arformoterol used to assess the plasma protein binding were higher than those achieved in plasma following inhalation of multiple doses of 50 mcg arformoterol. Metabolism <i>In vitro</i> profiling studies in hepatocytes and liver microsomes have shown that arformoterol is primarily metabolized by direct conjugation (glucuronidation) and secondarily by O-demethylation. At least five human uridine diphosphoglucuronosyltransferase (UGT) isozymes catalyze arformoterol glucuronidation <i>in vitro</i> . Two cytochrome P450 isozymes (CYP2D6 and secondarily CYP2C19) catalyze the O-demethylation of arformoterol. Arformoterol did not inhibit CYP1A2, CYP2A6, CYP2C9/10, CYP2C19, CYP2D6, CYP2E1, CYP3A4/5, or CYP4A9/11 enzymes at >1,000-fold higher concentrations than the expected peak plasma concentrations following a therapeutic dose. Arformoterol was almost entirely metabolized following oral administration of 35 mcg of radiolabeled arformoterol in eight healthy subjects. Direct conjugation of arformoterol with glucuronic acid was the major metabolic pathway. Most of the drug-related material in plasma and urine was in the form of glucuronide or sulfate conjugates of arformoterol. O-Demethylation and conjugates of the O-demethyl metabolite were relatively minor metabolites accounting for less than 17% of the dose recovered in urine and feces. Elimination After administration of a single oral dose of radiolabeled arformoterol to eight healthy male subjects, 63% of the total radioactive dose was recovered in urine and 11% in feces within 48 hours. A total of 89% of the total radioactive dose was recovered within 14 days, with 67% in urine and 22% in feces. Approximately 1% of the dose was recovered as unchanged arformoterol in urine over 14 days. Renal clearance was 8.9 L/hr for unchanged arformoterol in these subjects. In COPD patients given 15 mcg inhaled arformoterol twice a day for 14 days, the mean terminal half-life of arformoterol was 26 hours.

BROVANA® (arformoterol tartrate) Inhalation Solution

15 mcg * /2 mL

*potency expressed as arformoterol



MEDICATION GUIDE AND PRESCRIBING INFORMATION

BROVANA® (arformoterol tartrate) Inhalation Solution

15 mcg * /2 mL

*potency expressed as arformoterol

MEDICATION GUIDE AND PRESCRIBING INFORMATION

BROVANA® (arformoterol tartrate) Inhalation Solution 15 mcg * /2 mL
*potency expressed as arformoterol
For oral inhalation only
WARNING: ASTHMA RELATED DEATH Long-acting beta₂-adrenergic agonists (LABA) increase the risk of asthma-related death. Data from a large placebo-controlled US study that compared the safety of another long-acting beta₂-adrenergic agonist (salmeterol) or placebo added to usual asthma therapy showed an increase in asthma-related deaths in patients receiving salmeterol. This finding with salmeterol is considered a class effect of LABA, including arformoterol, the active ingredient in BROVANA (see WARNINGS). The safety and efficacy of BROVANA in patients with asthma have not been established. All LABA, including BROVANA, are contraindicated in patients with asthma without use of a long-term asthma control medication (see CONTRAINDICATIONS).
DESCRIPTION BROVANA (arformoterol tartrate) Inhalation Solution is a sterile, clear, colorless, aqueous solution of the tartrate salt of arformoterol, the (R,R)-enantiomer of formoterol. Arformoterol is a selective beta ₂ -adrenergic bronchodilator. The chemical name for arformoterol tartrate is formamide, N-[2-hydroxy-5-[(1R)-1-hydroxy-2-[[[(1R)-2-(4-methoxyphenyl)-1-methylethyl] amino]ethyl] phenyl]-, (2R,3R)-2,3-dihydroxybutanedioate (1:1 salt), and its established structural formula is as follows: <div style="display: flex; justify-content: space-around; align-items: center;"><div style="text-align: center;"><p>Chemical structure of arformoterol, showing the (R,R)-enantiomer of formoterol.</p></div><div style="text-align: center;"><p>Chemical structure of tartaric acid, the counterion of the tartrate salt.</p></div></div>
The molecular weight of <i>arformoterol tartrate</i> is 494.5 g/mol, and its empirical formula is C ₁₉ H ₂₄ N ₂ O ₄ • C ₄ H ₆ O ₆ (1:1 salt). It is a white to off-white solid that is slightly soluble in water. Arformoterol tartrate is the United States Adopted Name (USAN) for (R,R)-formoterol L-tartrate. BROVANA is supplied as 2 mL of arformoterol tartrate solution packaged in 2.1 mL unit-dose, low-density polyethylene (LDPE) ready-to-use vials. Each ready-to-use vial contains 15 mcg of arformoterol (equivalent to 22 mcg of arformoterol tartrate) in a sterile, isotonic saline solution, pH-adjusted to 5.0 with citric acid and sodium citrate. BROVANA requires no dilution before administration by nebulization. Like all other nebulized treatments, the amount delivered to the lungs will depend upon patient factors, the nebulizer used, and compressor performance. Using the PARI LC PLUS® nebulizer (with mouthpiece) connected to a PARI DUARA-NEB® 3000 compressor under <i>in vitro</i> conditions, the mean delivered dose from the mouthpiece (% nominal) was approximately 4.1 mcg (27.6%) at a mean flow rate of 3.3 L/min. The mean nebulization time was 6 minutes or less. BROVANA should be administered from a standard jet nebulizer at adequate flow rates via face mask or mouthpiece (see Dosage and Administration). Patients should be carefully instructed on the correct use of this drug product (please refer to the accompanying Medication Guide).
CLINICAL PHARMACOLOGY Mechanism of Action Arformoterol, the (R,R)-enantiomer of formoterol, is a selective long-acting beta ₂ -adrenergic receptor agonist (beta ₂ -agonist) that has two-fold greater potency than racemic formoterol (which contains both the (S,S) and (R,R)-enantiomers). The (S,S)-enantiomer is about 1,000-fold less potent as a beta ₂ -agonist than the (R,R)-enantiomer. While it is recognized that beta ₂ -receptors are the predominant adrenergic receptors in bronchial smooth muscle and beta ₂ -receptors are the predominant receptors in the heart, data indicate that there are also beta ₂ -receptors in the human heart comprising 10% to 50% of the total beta-adrenergic receptors. The precise function of these receptors has not been established, but they raise the possibility that even highly selective beta ₂ -agonists may have cardiac effects. The pharmacologic effects of beta ₂ -adrenoceptor agonist drugs, including arformoterol, are at least in part attributable to stimulation of intracellular adenylyl cyclase, the enzyme that catalyzes the conversion of adenosine triphosphate (ATP) to cyclic-3',5'-adenosine monophosphate (cyclic AMP). Increased intracellular cyclic AMP levels cause relaxation of bronchial smooth muscle and inhibition of release of mediators of immediate hypersensitivity from cells, especially from mast cells. <i>In vitro</i> tests show that arformoterol is an inhibitor of the release of mast cell mediators, such as histamine and leukotrienes, from the human lung. Arformoterol also inhibits histamine-induced plasma albumin extravasation in anesthetized guinea pigs and inhibits allergen-induced eosinophil influx in dogs with airway hyper-responsiveness. The relevance of these <i>in vitro</i> and animal findings to humans is unknown. Animal Pharmacology In animal studies investigating its cardiovascular effects, arformoterol induced dose-dependent increases in heart rate and decreases in blood pressure consistent with its pharmacology as a beta-adrenergic agonist. In dogs, at systemic exposures higher than anticipated clinically, arformoterol also induced exaggerated pharmacologic effects of a beta-adrenergic agonist on cardiac function as measured by electrocardiogram (sinus tachycardia, atrial premature beats, ventricular escape beats, PVCs). Studies in laboratory animals (minipigs, rodents, and dogs) have demonstrated the occurrence of arrhythmias and sudden death (with histologic evidence of myocardial necrosis) when beta-agonists and methylxanthines are administered concurrently. The clinical significance of these findings is unknown.
Pharmacokinetics The pharmacokinetics (PK) of arformoterol have been investigated in healthy subjects, elderly subjects, renally and hepatically impaired subjects, and chronic obstructive pulmonary disease (COPD) patients following the nebulization of the recommended therapeutic dose and doses up to 96 mcg. Absorption In COPD patients administered 15 mcg arformoterol every 12 hours for 14 days, the mean steady-state peak (R,R)-formoterol plasma concentration (C _{max}) and systemic exposure (AUC _{0-12h}) were 4.3 pg/mL and 34.5 pg•hr/mL, respectively. The median steady-state peak (R,R)-formoterol plasma concentration time (t _{max}) was observed approximately one half hour after drug administration. Systemic exposure to (R,R)-formoterol increased linearly with dose in COPD patients following arformoterol doses of 5 mcg, 15 mcg, or 25 mcg twice daily for 2 weeks or 15 mcg, 25 mcg, or 50 mcg once daily for 2 weeks. In a crossover study in patients with COPD, when arformoterol 15 mcg inhalation solution and 12 and 24 mcg formoterol fumarate inhalation powder (Foradil® Aerolizer™) was administered twice daily for 2 weeks, the accumulation index was approximately 2.5 based on the plasma (R,R)-formoterol concentrations in all three treatments. At steady state, geometric means of systemic exposure (AUC _{0-12h}) to (R,R)-formoterol following 15 mcg of arformoterol inhalation solution and 12 mcg of formoterol fumarate inhalation powder were 39.33 pg•hr/mL and 33.93 pg•hr/mL, respectively (ratio 1.16; 90% CI 1.00, 1.35), while the geometric means of the C _{max} were 4.30 pg/mL and 4.75 pg/mL, respectively (ratio 0.91; 90% CI 0.76, 1.09). In a study in patients with asthma, treatment with arformoterol 50 mcg with pre- and post-treatment with activated charcoal resulted in a geometric mean decrease in (R,R)-formoterol AUC _{0-6h} by 27% and C _{max} by 23% as compared to treatment with arformoterol 50 mcg alone. This suggests that a substantial portion of systemic drug exposure is due to pulmonary absorption. Distribution The binding of arformoterol to human plasma proteins <i>in vitro</i> was 52-65% at concentrations of 0.25, 0.5 and 1.0 ng/mL of radiolabeled arformoterol. The concentrations of arformoterol used to assess the plasma protein binding were higher than those achieved in plasma following inhalation of multiple doses of 50 mcg arformoterol. Metabolism <i>In vitro</i> profiling studies in hepatocytes and liver microsomes have shown that arformoterol is primarily metabolized by direct conjugation (glucuronidation) and secondarily by O-demethylation. At least five human uridine diphosphoglucuronosyltransferase (UGT) isozymes catalyze arformoterol glucuronidation <i>in vitro</i> . Two cytochrome P450 isozymes (CYP2D6 and secondarily CYP2C19) catalyze the O-demethylation of arformoterol. Arformoterol did not inhibit CYP1A2, CYP2A6, CYP2C9/10, CYP2C19, CYP2D6, CYP2E1, CYP3A4/5, or CYP4A9/11 enzymes at >1,000-fold higher concentrations than the expected peak plasma concentrations following a therapeutic dose. Arformoterol was almost entirely metabolized following oral administration of 35 mcg of radiolabeled arformoterol in eight healthy subjects. Direct conjugation of arformoterol with glucuronic acid was the major metabolic pathway. Most of the drug-related material in plasma and urine was in the form of glucuronide or sulfate conjugates of arformoterol. O-Demethylation and conjugates of the O-demethyl metabolite were relatively minor metabolites accounting for less than 17% of the dose recovered in urine and feces. Elimination After administration of a single oral dose of radiolabeled arformoterol to eight healthy male subjects, 63% of the total radioactive dose was recovered in urine and 11% in feces within 48 hours. A total of 89% of the total radioactive dose was recovered within 14 days, with 67% in urine and 22% in feces. Approximately 1% of the dose was recovered as unchanged arformoterol in urine over 14 days. Renal clearance was 8.9 L/hr for unchanged arformoterol in these subjects. In COPD patients given 15 mcg inhaled arformoterol twice a day for 14 days, the mean terminal half-life of arformoterol was 26 hours.

Special Populations

Gender

A population PK analysis indicated that there was no effect of gender upon the pharmacokinetics of arformoterol.

Race

The influence of race on arformoterol pharmacokinetics was assessed using a population PK analysis and data from healthy subjects. There was no clinically significant impact of race upon the pharmacokinetic profile of arformoterol.

Geriatric

The pharmacokinetic profile of arformoterol in 24 elderly subjects (aged 65 years or older) was compared to a younger cohort of 24 subjects (18-45 years) that were matched for body weight and gender. No significant differences in systemic exposure (AUC and C_{max}) were observed when the two groups were compared.

Pediatric

The pharmacokinetics of arformoterol have not been studied in pediatric subjects.

Hepatic Impairment

The pharmacokinetic profile of arformoterol was assessed in 24 subjects with mild, moderate, and severe hepatic impairment. The systemic exposure (C_{max} and AUC) to arformoterol increased 1.3 to 2.4-fold in subjects with hepatic impairment compared to 16 demographically matched healthy control subjects. No clear relationship between drug exposure and the severity of hepatic impairment was observed. BROVANA should be used cautiously in patients with hepatic impairment.

Renal Impairment

The impact of renal disease upon the pharmacokinetics of arformoterol was studied in 24 subjects with mild, moderate, or severe renal impairment. Systemic exposure (AUC and C_{max}) to arformoterol was similar in renally impaired patients compared with demographically matched healthy control subjects.

Pharmacogenetics

Arformoterol is eliminated through the action of multiple drug metabolizing enzymes. Direct glucuronidation of arformoterol is mediated by several UGT enzymes and is the primary elimination route. O-Desmethylation is a secondary route catalyzed by the CYP enzymes CYP2D6 and CYP2C19. In otherwise healthy subjects with reduced CYP2D6 and/or UGT1A1 enzyme activity, there was no impact on systemic exposure to arformoterol compared to subjects with normal CYP2D6 and/or UGT1A1 enzyme activities.

Pharmacodynamics

Systemic Safety and Pharmacokinetic/ Pharmacodynamic Relationships

The predominant adverse effects of inhaled beta₂-agonists occur as a result of excessive activation of systemic beta-adrenergic receptors. The most common adverse effects may include skeletal muscle tremor and cramps, insomnia, tachycardia, decreases in plasma potassium, and increases in plasma glucose.

Effects on Serum Potassium and Serum Glucose Levels

Changes in serum potassium and serum glucose were evaluated in a dose ranging study of twice daily (5 mcg, 15 mcg, or 25 mcg; 215 patients with COPD) and once daily (15 mcg, 25 mcg, or 50 mcg; 191 patients with COPD) BROVANA in COPD patients. At 2 and 6 hours post dose at week 0 (after the first dose), mean changes in serum potassium ranging from 0 to –0.3 mEq/L were observed in the BROVANA groups with similar changes observed after 2 weeks of treatment. Changes in mean serum glucose levels, ranging from a decrease of 1.2 mg/dL to an increase of 32.8 mg/dL were observed for BROVANA dose groups at both 2 and 6 hours post dose, both after the first dose and 14 days of daily treatment.

Electrophysiology

The effect of BROVANA on QT interval was evaluated in a dose ranging study following multiple doses of BROVANA 5 mcg, 15 mcg, or 25 mcg twice daily or 15 mcg, 25 mcg, or 50 mcg once daily for 2 weeks in patients with COPD. ECG assessments were performed at baseline, time of peak plasma concentration and throughout the dosing interval. Different methods of correcting for heart rate were employed, including a subject-specific method and the Fridericia method.

Relative to placebo, the mean change in subject-specific QT_c averaged over the dosing interval ranged from -1.8 to 2.7 msec, indicating little effect of BROVANA on cardiac repolarization after 2 weeks of treatment. The maximum mean change in subject-specific QT_c for the BROVANA 15 mcg twice daily dose was 17.3 msec, compared with 15.4 msec in the placebo group. No apparent correlation of QT_c with arformoterol plasma concentration was observed.

Electrocardiographic Monitoring in Patients with COPD

The effect of different doses of BROVANA on cardiac rhythm was assessed using 24-hour Holter monitoring in two 12-week double-blind, placebo-controlled studies of 1,456 patients with COPD (873 received BROVANA at 15 or 25 mcg twice daily or 50 mcg once daily doses; 293 received placebo; 290 received salmeterol). The 24-hour Holter monitoring occurred once at baseline, and up to 3 times during the 12-week treatment period. The rates of new-onset cardiac arrhythmias not present at baseline over the double-blind 12-week treatment period were similar (approximately 33-34%) for patients who received BROVANA 15 mcg twice daily to those who received placebo. There was a dose-related increase in new, treatment emergent arrhythmias seen in patients who received BROVANA 25 mcg twice daily and 50 mcg once daily, 37.6% and 40.1 %, respectively. The frequencies of new treatment emergent events of non-sustained (3-10 beat run) and sustained (>10 beat run) ventricular tachycardia were 7.4% and 1.1% in BROVANA 15 mcg twice daily and 6.9% and 1.0% in placebo. In patients who received BROVANA 25 mcg twice daily and 50 mcg once daily the frequencies of non-sustained (6.2% and 8.2%, respectively) and sustained ventricular tachycardia (1.0% and 1.0%, respectively) were similar. Five cases of ventricular tachycardia were reported as adverse events (1 in BROVANA 15 mcg twice daily and 4 in placebo), with two of these events leading to discontinuation of treatment (2 in placebo). There were no baseline occurrences of atrial fibrillation/ flutter observed on 24-hour Holter monitoring in patients treated with BROVANA 15 mcg twice daily or placebo. New, treatment emergent atrial fibrillation/ flutter occurred in 0.4% of patients who received BROVANA 15 mcg twice daily and 0.3% of patients who received placebo. There was a dose-related increase in the frequency of atrial fibrillation/ flutter reported in the BROVANA 25 mcg twice daily and 50 mcg once daily dose groups of 0.7% and 1.4%, respectively. Two cases of atrial fibrillation/ flutter were reported as adverse events (1 in BROVANA 15 mcg twice daily and 1 in placebo).


Dose-related increases in mean maximum change in heart rate in the 12 hours after dosing were also observed following 12 weeks of dosing with BROVANA 15 mcg twice daily (8.8 bpm), 25 mcg twice daily (9.9 bpm) and 50 mcg once daily (12 bpm) versus placebo (8.5 bpm).
Tachyphylaxis/Tolerance
In two placebo-controlled clinical trials in patients with COPD involving approximately 725 patients in each, the overall efficacy of BROVANA was maintained throughout the 12-week trial duration. However, tolerance to the bronchodilator effect of BROVANA was observed after 6 weeks of dosing, evidenced by a decrease in bronchodilator effect as measured by FEV₁. FEV₁ improvement at the end of the 12-hour dosing interval decreased by approximately one third (22.1% mean improvement after the first dose compared to 14.6% at week 12). Tolerance to the FEV₁ bronchodilator effect of BROVANA was not accompanied by other clinical manifestations of tolerance in these trials.

CLINICAL TRIALS

Adult COPD Trials

BROVANA (arformoterol tartrate) Inhalation Solution was studied in two identical, 12 week, double-blind, placebo- and active-controlled, randomized, multi-center, parallel group trials conducted in the United States (Clinical Trial A and Clinical Trial B). A total of 1,456 adult patients (age range: 34 to 89 years; mean age: 63 years) with COPD who had a mean FEV₁ of 1.3 L (42% of predicted) were enrolled in the two clinical trials. The diagnosis of COPD was based on a prior clinical diagnosis of COPD, a smoking history (greater than 15 pack-years), age (at least 35 years), spirometry results (baseline FEV₁ ≤65% of predicted value and ≤0.70 L, and a FEV₁/forced vital capacity (FVC) ratio ≤70%). About 80% of patients in these studies had bronchodilator reversibility, defined as a 10% or greater increase in FEV₁ after inhalation of 2 actuations (180 mcg racemic albuterol from a metered dose inhaler). Both trials compared BROVANA 15 mcg twice daily (288 patients), 25 mcg twice daily (292 patients), 50 mcg once daily (293 patients) with placebo (293 subjects). Both trials included salmeterol inhalation aerosol, 42 mcg twice daily as an active comparator (290 patients).

In both 12-week trials, BROVANA 15 mcg twice daily resulted in significantly greater post-dose bronchodilation (as measured by percent change from study baseline FEV₁ at the end of the dosing interval over the 12 weeks of treatment, the primary efficacy endpoint) compared to placebo. Compared to BROVANA 15 mcg twice daily, BROVANA 25 mcg twice daily and 50 mcg once daily did not provide sufficient additional benefit on a variety of endpoints, including FEV₁, to support the use of higher doses. Plots of the mean change in FEV₁ values obtained over the 12 hours after dosing for the BROVANA 15 mcg twice daily dose group and for the placebo group are provided in Figures 1 and 2 for Clinical Trial A, below. The plots include mean FEV₁ change observed after the first dose and after 12 weeks of treatment. The results from Clinical Trial B were similar.



MEDICATION GUIDE

BROVANA® (*Brô vâ`-nah*) (arformoterol tartrate) Inhalation Solution

BROVANA is only for use with a nebulizer.

Read the Medication Guide that comes with BROVANA before you start using it and each time you get a refill. There may be new information. This Medication Guide does not take the place of talking to your healthcare provider about your medical condition or treatment.

What is the most important information I should know about BROVANA?

BROVANA can cause serious side effects, including:

- People with asthma, who take long-acting beta₂ adrenergic agonist (LABA) medicines, such as BROVANA, have an increased risk of death from asthma problems.**
- It is not known if LABA medicines, such as BROVANA, increase the risk of death in people with chronic obstructive pulmonary disease (COPD).
- Get emergency medical care if:**
 - breathing problems worsen quickly**
 - you use your rescue inhaler medicine, but it does not relieve your breathing problems**

What is BROVANA?

BROVANA is used long term, 2 times each day (morning and evening), in controlling symptoms of chronic obstructive pulmonary disease (COPD) in adults with COPD.

BROVANA is only for use with a nebulizer.

LABA medicines such as BROVANA help the muscles around the airways in your lungs stay relaxed to prevent symptoms, such as wheezing, cough, chest tightness, and shortness of breath.

BROVANA should not be used in children. **It is not known if BROVANA is safe and effective in children.**

It is not known if BROVANA is safe and effective in people with asthma.

Who should not use BROVANA?

Do not use BROVANA if you:

- have had a serious allergic reaction to arformoterol, formoterol, or any of the ingredients in BROVANA. Ask your healthcare provider if you are not sure. See the end of this Medication Guide for a complete list of ingredients in BROVANA.
- have asthma without using a long-term asthma control medicine.

What should I tell my healthcare provider before using BROVANA?

Tell your healthcare provider about all of your health conditions, including if you:

- have heart problems
- have high blood pressure
- have seizures
- have thyroid problems
- have diabetes
- have liver problems
- are pregnant or planning to become pregnant. It is not known if BROVANA can harm your unborn baby.
- are breastfeeding. It is not known if BROVANA passes into your milk and if it can harm your baby.

Tell your healthcare provider about all the medicines you take including prescription and non-prescription medicines, vitamins and herbal supplements. BROVANA and certain other medicines may interact with each other. This may cause serious side effects.

Know the medicines you take. Keep a

- **nervousness**
- **leg cramps**
- **high blood potassium**
- **shortness of breath**
- **rash**
- **fever**
- **increased white blood cells**
- **vomiting**
- **tiredness**
- **leg swelling**
- **chest congestion or bronchitis**

Tell your healthcare provider if you get any side effect that bothers you or that does not go away.

These are not all the side effects with BROVANA. Ask your healthcare provider or pharmacist for more information. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store BROVANA?

- Store BROVANA in a refrigerator between 36° to 46°F (2° to 8°C) in the protective foil pouch. Protect from light and excessive heat. **Do not open a sealed pouch until you are ready to use a dose of BROVANA. After opening the pouch, unused ready-to-use vials should be returned to, and stored in, the pouch. An opened ready-to-use vial should be used right away.** BROVANA may be used directly from the refrigerator.
- BROVANA may also be stored at room temperature between 68°F to 77°F (20°C to 25°C) for up to 6 weeks (42 days). If stored at room temperature, discard BROVANA if it is not used after 6 weeks or if past the expiration date, whichever is sooner. Space is provided on the packaging to record room temperature storage times.
- Do not use BROVANA after the expiration date provided on the foil pouch and ready-to-use vial.
- BROVANA should be colorless. Discard BROVANA if it is not colorless.
- **Keep BROVANA and all medicines out of the reach of children.**

General Information about BROVANA

Medicines are sometimes prescribed for purposes not mentioned in a Medication Guide. Do not use BROVANA for a condition for which it was not prescribed. Do not give BROVANA to other people, even if they have the same condition. It may harm them.

This Medication Guide summarizes the most important information about BROVANA. If you would like more information, talk with your healthcare provider. You can ask your healthcare provider or pharmacist for information about BROVANA that was written for healthcare professionals.

- For customer service, call 1-888-394-7377.
- To report side effects, call 1-877-737-7226.
- For medical information, call 1-800-739-0565.

Instructions for Using BROVANA (arformoterol tartrate) Inhalation Solution

BROVANA is used only in a standard jet nebulizer machine connected to an air compressor. Make sure you know how to use your nebulizer machine before you use it to breathe-in BROVANA or other medicines.

Do not mix BROVANA with other medicines in your nebulizer machine.

BROVANA comes sealed in a foil pouch. Do not open a sealed pouch until you are ready to use a dose of BROVANA. After opening the pouch, unused ready-to-use vials should be returned to, and stored in, the pouch. An opened ready-to-use vial should be used right away.

1. Open the foil pouch by tearing on the rough edge along the seam of the pouch. Remove a ready-to-use vial of BROVANA.

2. Carefully twist open the top of the ready-to-use vial and use it right away (**Figure 1**).




Figure 1

3. Squeeze all of the medicine from the ready-to-use vial into the nebulizer medicine cup (reservoir) (**Figure 2**).




Figure 2

4. Connect the nebulizer reservoir to the mouthpiece (**Figure 3**) or face mask (**Figure 4**).



Figure 3



Figure 4

5. Connect the nebulizer to the compressor (**Figure 5**).




Figure 5

6. Sit in a comfortable, upright position. Place the mouthpiece in your mouth (**Figure 6**) (or put on the face mask) and turn on the compressor.




Figure 6

7. Breathe as calmly, deeply, and evenly as possible until no more mist is formed in the nebulizer reservoir. It takes about 5 to 10 minutes for each treatment.

8. Clean the nebulizer (see manufacturer's instructions).

Rx Only

This Medication Guide has been approved by the Food and Drug Administration.




Manufactured for:
Sunovion Pharmaceuticals Inc.
Marlborough, MA 01752 USA

BROVANA is a registered trademark of Sunovion Pharmaceuticals Inc.

January 2011
901005R06

Figure 2 Mean Change in FEV₁ Over Time for Clinical Trial A at Week 12



BROVANA 15 mcg twice daily significantly improved bronchodilation compared to placebo over the 12 hours after dosing (FEV₁, AUC_{0-12h}). This improvement was maintained over the 12 week study period.

Following the first dose of BROVANA 15 mcg, the median time to onset of bronchodilation, defined by an FEV₁ increase of 15%, occurred at 6.7 min. When defined as an increase in FEV₁ of 12% and 200 mL, the time to onset of bronchodilation was 20 min after dosing. Peak bronchodilator effect was generally seen within 1-3 hours of dosing.

In both clinical trials, compared to placebo, patients treated with BROVANA demonstrated improvements in peak expiratory flow rates, supplemental ipratropium and rescue albuterol use.

INDICATIONS AND USAGE

BROVANA (arformoterol tartrate) Inhalation Solution is indicated for the long term, twice daily (morning and evening) maintenance treatment of bronchoconstriction in patients with chronic obstructive pulmonary disease (COPD), including chronic bronchitis and emphysema. BROVANA is for use by nebulization only.

CONTRAINDICATIONS

BROVANA (arformoterol tartrate) Inhalation Solution is contraindicated in patients with a history of hypersensitivity to arformoterol, racemic formoterol or to any other components of this product.

All LABA, including BROVANA, are contraindicated in patients with asthma without use of a long-term asthma control medication (see **WARNINGS**).

WARNINGS

• ASTHMA RELATED DEATH

Long-acting beta₂-adrenergic agonists increase the risk of asthma-related death. The safety and efficacy of BROVANA in patients with asthma have not been established. All LABA, including BROVANA, are contraindicated in patients with asthma without use of a long-term asthma control medication (see CONTRAINDICATIONS).

• A 28-week, placebo-controlled US study comparing the safety of salmeterol with placebo, each added to usual asthma therapy, showed an increase in asthma-related deaths in patients receiving salmeterol (13/13,176 in patients treated with salmeterol vs. 3/13,179 in patients treated with placebo; RR 4.37, 95% CI 1.25, 15.34). The increased risk of asthma-related death is considered a class effect of the long-acting beta₂-adrenergic agonists, including BROVANA. No study adequate to determine whether the rate of asthma related death is increased in patients treated with BROVANA has been conducted.

• Clinical studies with racemic formoterol (Foradil® Aerolizer™) suggested a higher incidence of serious asthma exacerbations in patients who received racemic formoterol than in those who received placebo. The sizes of these studies were not adequate to precisely quantify the differences in serious asthma exacerbation rates between treatment groups.

• **The studies described above enrolled patients with asthma. Data are not available to determine whether the rate of death in patients with COPD is increased by long-acting beta₂-adrenergic agonists.**

• **BROVANA is indicated for the long term, twice daily (morning and evening) maintenance treatment for bronchoconstriction in chronic obstructive pulmonary disease (COPD), and is not indicated for the treatment of acute episodes of bronchospasm, i.e., rescue therapy.**

• **BROVANA should not be initiated in patients with acutely deteriorating COPD, which may be a life-threatening condition. The use of BROVANA in this setting is inappropriate.**

• **BROVANA should not be used in children as the safety and efficacy of BROVANA have not been established in pediatric patients.**

• **BROVANA should not be used in conjunction with other inhaled, long-acting beta₂-agonists. BROVANA should not be used with other medications containing long-acting beta₂-agonists.**

• **When beginning treatment with BROVANA, patients who have been taking inhaled, short-acting beta₂-agonists on a regular basis (e.g., four times a day) should be instructed to discontinue the regular use of these drugs and use them only for symptomatic relief of acute respiratory symptoms.**

• See **PRECAUTIONS, Information for Patients and the accompanying Medication Guide.**

Paradoxical Bronchospasm

As with other inhaled beta₂-agonists, BROVANA can produce paradoxical bronchospasm that may be life-threatening. If paradoxical bronchospasm occurs, BROVANA should be discontinued immediately and alternative therapy instituted.

Deterioration of Disease

COPD may deteriorate acutely over a period of hours or chronically over several days or longer. If BROVANA no longer controls the symptoms of bronchoconstriction, or the patient's inhaled, short-acting beta₂-agonist becomes less effective or the patient needs more inhalation of short-acting beta₂-agonist than usual, these may be markers of deterioration of disease. In this setting, a re-evaluation of the patient and the COPD treatment regimen should be undertaken at once. Increasing the daily dosage of BROVANA beyond the recommended 15 mcg twice daily dose is not appropriate in this situation.

Cardiovascular Effects

BROVANA, like other beta₂-agonists, can produce a clinically significant cardiovascular effect in some patients as measured by increases in pulse rate, blood pressure, and/or symptoms. Although such effects are uncommon after administration of BROVANA at the recommended dose, if they occur, the drug may need to be discontinued. In addition, beta₂-agonists have been reported to produce ECG changes, such as flattening of the T wave, prolongation of the QT interval, and ST segment depression. The clinical significance of these findings is unknown. BROVANA, as with other sympathomimetic amines, should be used with caution in patients with cardiovascular disorders, especially coronary insufficiency, cardiac arrhythmias, and hypertension (see **PRECAUTIONS, General**).

Immediate Hypersensitivity Reactions

Immediate hypersensitivity reactions may occur after administration of BROVANA as demonstrated by cases of anaphylactic reaction, urticaria, angioedema, rash and bronchospasm.

Do Not Exceed Recommended Dose

Fatalities have been reported in association with excessive use of inhaled sympathomimetic drugs. As with other inhaled beta₂-adrenergic drugs, BROVANA should not be used more often, at higher doses than recommended, or with other long-acting beta-agonists.

PRECAUTIONS

General

BROVANA (arformoterol tartrate) Inhalation Solution should not be used to treat acute symptoms of COPD. BROVANA has not been studied in the relief of acute symptoms and extra doses should not be used for that purpose. When prescribing BROVANA, the physician should also provide the patient with an inhaled, short-acting beta₂-agonist for treatment of COPD symptoms that occur acutely, despite regular twice-daily (morning and evening) use of BROVANA. Patients should also be cautioned that increasing inhaled beta₂-agonist use is a signal of deteriorating disease for which prompt medical attention is indicated (see **Information for Patients** and the accompanying **Medication Guide**).

BROVANA, like other sympathomimetic amines, should be used with caution in patients with cardiovascular disorders, especially coronary insufficiency, cardiac arrhythmias, and hypertension; in patients with convulsive disorders or thyrotoxicosis; and in patients who are unusually responsive to sympathomimetic amines. Clinically significant changes in systolic and/or diastolic blood pressure, pulse rate and electrocardiograms have been seen infrequently in individual patients in controlled clinical studies with arformoterol tartrate. Doses of the related beta₂-agonist albuterol, when administered intravenously, have been reported to aggravate preexisting diabetes mellitus and ketoacidosis.

Beta-agonist medications may produce significant hypokalemia in some patients, possibly through intracellular shunting, which has the potential to produce adverse cardiovascular effects. The decrease in serum potassium is usually transient, not requiring supplementation.

Clinically significant changes in blood glucose and/or serum potassium were infrequent during clinical studies with long-term administration of BROVANA at the recommended dose.

Information for Patients

Patients should be instructed to read the accompanying Medication Guide with each new prescription and refill. The complete text of the Medication Guide is reprinted at the end of this document. Patients should be given the following information:

1. Patients should be informed that long-acting beta₂-adrenergic agonists, such as BROVANA, increase the risk of asthma-related death. All LABA, including BROVANA, should not be used in patients with asthma without use of a long-term asthma control medication (see **CONTRAINDICATIONS**).

2. BROVANA is not indicated to relieve acute respiratory symptoms and extra doses should not be used for that purpose. Acute symptoms should be treated with an inhaled, short-acting, beta₂-agonist (the health-care provider should prescribe the patient with such medication and instruct the patient in how it should be used). Patients should be instructed to seek medical attention if their symptoms worsen, if BROVANA treatment becomes less effective, or if they need more inhalations of a short-acting beta₂-agonist than usual. Patients should not inhale more than one dose at any one time. The daily dosage of BROVANA should not exceed one ready-to-use vial (15 mcg) by inhalation twice daily (30 mcg total daily dose).
3. Patients should be informed that treatment with beta₂-agonists may lead to adverse events which include palpitations, chest pain, rapid heart rate, tremor, or nervousness.
4. Patients should be instructed to use BROVANA by nebulizer only and not to inject or swallow this inhalation solution.
5. Patients should protect BROVANA ready-to-use vials from light and excessive heat. The protective foil pouches should be stored under refrigeration between 2°C and 8°C (36°–46°F). They should not be used after the expiration date stamped on the container. After opening the pouch, unused ready-to-use vials should be returned to, and stored in, the pouch. An opened ready-to-use vial should be used right away. Discard any ready-to-use vial if the solution is not colorless.
6. The drug compatibility (physical and chemical), efficacy and safety of BROVANA when mixed with other drugs in a nebulizer have not been established.
7. Women should be advised to contact their physician if they become pregnant or if they are nursing.
8. It is important that patients understand how to use BROVANA appropriately and how it should be used in relation to other medications to treat COPD they are taking (see the accompanying **Medication Guide** and the **Instructions for Using BROVANA**).

Drug Interactions

If additional adrenergic drugs are to be administered by any route, they should be used with caution because the pharmacologically predictable sympathetic effects of BROVANA may be potentiated.

When paroxetine, a potent inhibitor of CYP2D6, was co-administered with BROVANA at steady-state, exposure to either drug was not altered. Dosage adjustments of BROVANA are not necessary when the drug is given concomitantly with potent CYP2D6 inhibitors.

Concomitant treatment with methylxanthines (aminophylline, theophylline), steroids, or diuretics may potentiate any hypokalemic effect of adrenergic agonists.

The ECG changes and/or hypokalemia that may result from the administration of non-potassium sparing diuretics (such as loop or thiazide diuretics) can be acutely worsened by beta-agonists, especially when the recommended dose of the beta-agonist is exceeded. Although the clinical significance of these effects is not known, caution is advised in the co-administration of beta-agonists with non-potassium sparing diuretics.

BROVANA, as with other beta₂-agonists, should be administered with extreme caution to patients being treated with monoamine oxidase inhibitors, tricyclic antidepressants, or drugs known to prolong the QTc interval because the action of adrenergic agonists on the cardiovascular system may be potentiated by these agents. Drugs that are known to prolong the QTc interval have an increased risk of ventricular arrhythmias. The concurrent use of intravenously or orally administered methylxanthines (e.g., aminophylline, theophylline) by patients receiving BROVANA has not been completely evaluated. In two combined 12-week placebo controlled trials that included BROVANA doses of 15 mcg twice daily, 25 mcg twice daily, and 50 mcg once daily, 54 of 873 BROVANA-treated subjects received concomitant theophylline at study entry. In a 12-month controlled trial that included a 50 mcg once daily BROVANA dose, 30 of the 528 BROVANA-treated subjects received concomitant theophylline at study entry. In these trials, heart rate and systolic blood pressure were approximately 2-3 bpm and 6-8 mm Hg higher, respectively, in subjects on concomitant theophylline compared with the overall population.

Beta-adrenergic receptor antagonists (beta-blockers) and BROVANA may interfere with the effect of each other when administered concurrently. Beta-blockers not only block the therapeutic effects of beta-agonists, but may produce severe bronchospasm in COPD patients. Therefore, patients with COPD should not normally be treated with beta-blockers. However, under certain circumstances, e.g., as prophylaxis after myocardial infarction, there may be no acceptable alternatives to the use of beta-blockers in patients with COPD. In this setting, cardioselective beta-blockers could be considered, although they should be administered with caution.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term studies were conducted in mice using oral administration and rats using inhalation administration to evaluate the carcinogenic potential of arformoterol.

In a 24-month carcinogenicity study in CD-1 mice, arformoterol caused a dose-related increase in the incidence of uterine and cervical endometrial stromal polyps and stromal cell sarcoma in female mice at oral doses of 1 mg/kg and above (AUC exposure approximately 70 times adult exposure at the maximum recommended daily inhalation dose).

In a 24-month carcinogenicity study in Sprague-Dawley rats, arformoterol caused a statistically significant increase in the incidence of thyroid gland c-cell adenoma and carcinoma in female rats at an inhalation dose of 200 mcg/kg (AUC exposure approximately 130 times adult exposure at the maximum recommended daily inhalation dose). There were no tumor findings with an inhalation dose of 40 mcg/kg (AUC exposure approximately 55 times adult exposure at the maximum recommended daily inhalation dose).

Arformoterol was not mutagenic or clastogenic in the following tests: mutagenicity tests in bacteria, chromosome aberration analyses in mammalian cells, and micronucleus test in mice.

Arformoterol had no effects on fertility and reproductive performance in rats at oral doses up to 10 mg/kg (approximately 2700 times the maximum recommended daily inhalation dose in adults on a mg/m² basis).

Pregnancy; Teratogenic Effects

Pregnancy Category C

Arformoterol has been shown to be teratogenic in rats based upon findings of omphalocele (umbilical hernia), a malformation, at oral doses of 1 mg/kg and above (AUC exposure approximately 370 times adult exposure at the maximum recommended daily inhalation dose). Increased pup loss at birth and during lactation and decreased pup weights were observed in rats at oral doses of 5 mg/kg and above (AUC exposure approximately 1100 times adult exposure at the maximum recommended daily inhalation dose). Delays in development were evident with an oral dose of 10 mg/kg (AUC exposure approximately 2400 times adult exposure at the maximum recommended daily inhalation dose).

Arformoterol has been shown to be teratogenic in rabbits based upon findings of malpositioned right kidney, a malformation, at oral doses of 20 mg/kg and above (AUC exposure approximately 8400 times adult exposure at the maximum recommended daily inhalation dose). Malformations including brachydactyly, bulbous aorta, and liver cysts were observed at doses of 40 mg/kg and above (approximately 22,000 times the maximum recommended daily inhalation dose in adults on a mg/m² basis). Malformation including adactyly, lobular dysgenesis of the lung, and interventricular septal defect were observed at 80 mg/kg (approximately 43,000 times the maximum recommended daily inhalation dose in adults on a mg/m² basis). Embryolethality was observed at 80 mg/kg/day (approximately 43,000 times the maximum recommended daily inhalation dose in adults on a mg/m² basis). Decreased pup body weights were observed at doses of 40 mg/kg/day and above (approximately 22,000 times the maximum recommended daily inhalation dose in adults on a mg/m² basis). There were no teratogenic findings in rabbits with oral dose of 10 mg/kg and lower (AUC exposure approximately 4900 times adult exposure at the maximum recommended daily inhalation dose).

There are no adequate and well-controlled studies in pregnant women. BROVANA should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Use in Labor and Delivery

There are no human studies that have investigated the effects of BROVANA on preterm labor or labor at term.

Because beta-agonists may potentially interfere with uterine contractility, BROVANA should be used during labor and delivery only if the potential benefit justifies the potential risk.

Nursing Mothers

In reproductive studies in rats, arformoterol was excreted in the milk. It is not known whether arformoterol is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when BROVANA is administered to a nursing woman.

Pediatric

BROVANA is approved for use in the long term maintenance treatment of bronchoconstriction associated with chronic obstructive pulmonary disease, including chronic bronchitis and emphysema. This disease does not occur in children. The safety and effectiveness of BROVANA in pediatric patients have not been established.

Geriatric

Of the 873 patients who received BROVANA in two placebo-controlled clinical studies in adults with COPD, 391 (45%) were 65 years of age or older while 96 (11%) were 75 years of age or older. No overall differences in safety or effectiveness were observed between these subjects and younger subjects. Among subjects age 65 years and older, 129 (33%) received BROVANA at the recommended dose of 15 mcg twice daily, while the remainder received higher doses. ECG alerts for ventricular ectopy in patients 65 to <75 years of age were comparable among patients receiving 15 mcg twice daily, 25 mcg twice daily, and placebo (3.9%, 5.2%, and 7.1%, respectively). A higher frequency (12.4%) was observed when BROVANA was dosed at 50 mcg once daily. The clinical significance of this finding is not known. Other reported clinical experience has not identified differences in responses between the elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

ADVERSE REACTIONS

Experience in Adult Patients with COPD

Of the 1,456 COPD patients in the two 12-week, placebo-controlled trials, 288 were treated with BROVANA (arformoterol tartrate) Inhalation Solution 15 mcg twice daily and 293 were treated with placebo. Doses of 25 mcg twice daily and 50 mcg once daily were also evaluated. The numbers and percent of patients who reported adverse events were comparable in the 15 mcg twice daily and placebo groups.

The following table shows adverse events where the frequency was greater than or equal to 2% in the BROVANA 15 mcg twice daily group and where the rates of adverse events in the BROVANA 15 mcg twice daily group exceeded placebo. Ten adverse events demonstrated a dose relationship: asthenia, fever, bronchitis, COPD, headache, vomiting, hyperkalemia, leukocytosis, nervousness, and tremor.

Table 1: Number of Patients Experiencing Adverse Events from Two 12-Week, Double-Blind, Placebo Controlled Clinical Trials

	BROVANA 15 mcg twice daily		Placebo	
	n	(%)	n	(%)
Total Patients	288	(100)	293	(100)
Pain	23	(8)	16	(5)
Chest Pain	19	(7)	19	(6)
Back Pain	16	(6)	6	(2)
Diarrhea	16	(6)	13	(4)
Sinusitis	13	(5)	11	(4)
Leg Cramps	12	(4)	6	(2)
Dyspnea	11	(4)	7	(2)
Rash	11	(4)	5	(2)
Flu Syndrome	10	(3)	4	(1)
Peripheral Edema	8	(3)	7	(2)
Lung Disorder*	7	(2)	2	(1)

*Reported terms coded to "Lung Disorder" were predominantly pulmonary or chest congestion.

Adverse events occurring in patients treated with BROVANA 15 mcg twice daily with a frequency of <2%, but greater than placebo were as follows:

Body as a Whole: abscess, allergic reaction, digitalis intoxication, fever, hernia, injection site pain, neck rigidity, neoplasm, pelvic pain, retroperitoneal hemorrhage

Cardiovascular: arteriosclerosis, atrial flutter, AV block, congestive heart failure, heart block, myocardial infarct, QT interval prolonged, supraventricular tachycardia, inverted T-wave

Digestive: constipation, gastritis, melena, oral moniliasis, periodontal abscess, rectal hemorrhage

Metabolic and Nutritional Disorders: dehydration, edema, glucose tolerance decreased, gout, hyperglycemia, hyperlipidemia, hypoglycemia, hypokalemia

Musculoskeletal: arthralgia, arthritis, bone disorder, rheumatoid arthritis, tendinous contracture

Nervous: agitation, cerebral infarct, circumoral paresthesia, hypokinesia, paralysis, somnolence, tremor

Respiratory: carcinoma of the lung, respiratory disorder, voice alteration